

Kinetics of electrophysical properties of diluted aqueous colloidal solutions of fullerene $C_{60}@H_2O_n$

**M. A. Vinnikov¹, M. T. Pohribnyi¹, O. V. Dolbyn¹,
R. M. Basnukaeva¹, L. M. Buravtseva¹,
S. V. Cherednychenko¹**

¹*B. Verkin Institute for Low Temperature Physics and Engineering of the NAS of Ukraine,
47 Nauky Ave., Kharkiv, 61103, Ukraine*

e-mail: vinnikov@ilt.kharkov.ua

$C_{60}@H_2O_n$ mixture




The solid phase of the $C_{60}@H_2O_n$ mixture




Colloidal aqueous solution of the $C_{60}@H_2O_n$ mixture


Frequency dependences of the dielectric loss tangent δ and dielectric constant ϵ of aqueous colloidal solutions of fullerene $C_{60}@H_2O_n$



Time dependences of the dielectric loss tangent δ of aqueous colloidal solutions of fullerene $C_{60}@H_2O_n$



Time dependences of the dielectric constant ϵ of aqueous colloidal solutions of fullerene $C_{60}@H_2O_n$



Acknowledgement

This work was partly supported by the

- National Academy of Sciences of Ukraine
[Contract no 2H/48-26]
- National Research Foundation of Ukraine
[Grant 2023.03/0012]



Thank you for your attention!